

8 Permafrost

8.1 Scenarios

The simulation scenarios for models only participating as permafrost models are described below. Assuming that for the relevant regions “other human influences” only play a minor role, i.e. the regional simulations can be done as “naturalized” runs (**nosoc**). Results from permafrost modules embedded in global biomes models should be reported for the biomes model simulations specified in Section 6 and the extension beyond 2299 described below. Since the pre-industrial simulations are an important part of the experiments, the spin-up has to finish before the pre-industrial simulations start. The spin-up should be using pre-industrial climate (**picontrol**) and year 1860 levels of “other human influences”. For this reason, the pre-industrial climate data should be replicated as often as required. The precise implementation of the spin up will be model specific, the description of which will be part of the reporting process.

IMPORTANT: Please contact the permafrost sector coordinators (see <https://www.isimip.org/about/#sectors-and-contacts>) before starting permafrost simulations.

Climate & CO ₂ scenarios	
picontrol	Pre-industrial climate and 286ppm CO ₂ concentration. The climate data for the entire period (1661-2299) are unique - no (or little) recycling of data has taken place.
historical	Historical climate and CO ₂ concentration.
rcp26	Future climate and CO ₂ concentration from RCP2.6.
rcp60	Future climate and CO ₂ concentration from RCP6.0.
rcp85	Future climate and CO ₂ concentration from RCP8.5.
2299rcp26	Repeating climate between 2270 and 2299 for additional 200 years up to 2500 (or equilibrium if possible), CO ₂ fixed at year 2299 levels.
2005co2	Fixed year 2005 CO ₂ concentration.
Human influence & land-use scenarios	
nosoc*	No human influences.

* Simulations using the human influence & land-use scenario design of the Biomes sector are also accepted (see Section 6.1), but nosoc is preferred.

Table 39 ISIMIP2b scenario specification for the permafrost simulations. Simulations using the scenario specifications of the Biomes sector are also accepted (see Section 6.1), but the permafrost specifications are preferred.

Experiment	Input	Pre-industrial	Historical	Future	Extended future	Beyond 2299
------------	-------	----------------	------------	--------	-----------------	-------------

		1661-1860	1861-2005	2006-2099	2100-2299		
I	no climate change, pre-industrial CO ₂	Climate & CO ₂	picontrol	not simulated	not simulated	not simulated	not simulated
	no other human influences	Human & LU	nosoc				
II	RCP2.6 climate & CO ₂	Climate & CO ₂	Experiment I	historical	rcp26	rcp26	2299rcp26
	no other human influences	Human & LU		nosoc	nosoc	nosoc	nosoc
Ila	RCP2.6 climate, CO ₂ varying until 2005, then fixed at 2005 levels thereafter	Climate & CO ₂	Experiment I	Experiment II	rcp26, 2005co2	rcp26, 2005co2	2299rcp26, 2005co2
	no other human influences	Human & LU			nosoc	nosoc	nosoc
III	RCP6.0 climate & CO ₂	Climate & CO ₂	Experiment I	Experiment II	rcp60	not simulated	not simulated
	no other human influences	Human & LU			nosoc		
IV-VII	Not simulated						
VIII	RCP8.5 climate & CO ₂	Climate & CO ₂	Experiment I	Experiment II	rcp85	not simulated	not simulated
	no other human influences	Human & LU			nosoc		

8.2 Output data

Table 40 Variables to be reported by permafrost models.

Variable name (long name)	Variable name	Unit (NetCDF format)	Resolution	Comment
---------------------------	---------------	----------------------	------------	---------

Essential outputs					
Temperature of Soil	tsl	K	per gridcell	daily (monthly)	Temperature of each soil layer. Reported as "missing" for grid cells occupied entirely by "sea". THIS IS THE MOST IMPORTANT VARIABLE. Also need depths in meters. Daily is preferred over monthly.
Pools (as Biomes output Table)					
Carbon Mass in Vegetation biomass	cveg-<pft>	kg m-2	per pft and gridcell total	annual	Gridcell total cveg is essential. Per PFT information is desirable.
Carbon Mass in aboveground vegetation biomass	cvegag-<pft>	kg m-2	per pft and gridcell total	annual	Gridcell total cvegag is essential. Per PFT information is desirable.
Carbon Mass in belowground vegetation biomass	cvebg-<pft>	kg m-2	per pft and gridcell total	annual	Gridcell total cvebg is essential. Per PFT information is desirable.
Carbon Mass in Litter Pool	clitter	kg m-2	per gridcell total	annual	Info for each individual pool.
Carbon Mass in Soil Pool	csoil	kg m-2	per gridcell total	annual	Info for each individual pool.
*Total Carbon Mass in Soil Pool	soilc	kg m-2	per gridcell total	annual	Integrated over the entire soil depth
Fluxes (as Biomes output Table)					
Carbon Mass Flux out of atmosphere due to Gross Primary Production on Land	gpp	kg m-2 s-1	gridcell total	daily (monthly)	
Carbon Mass Flux out of atmosphere due to Gross Primary Production on Land	gpp-<pft>	kg m-2 s-1	per pft	annual	
Carbon Mass Flux into atmosphere due to Autotrophic (Plant) Respiration on Land	ra	kg m-2 s-1	gridcell total	daily (monthly)	
Carbon Mass Flux out of atmosphere due to Net	npp	kg m-2 s-1	gridcell total	daily (monthly)	

Primary Production on Land					
Carbon Mass Flux out of atmosphere due to Net Primary Production on Land	npp-<pft>	kg m-2 s-1	per pft	annual	
Carbon Mass Flux into atmosphere due to Heterotrophic Respiration on Land	rh	kg m-2 s-1	gridcell total	daily (monthly)	
Carbon Mass Flux into atmosphere due to total Carbon emissions from Fire	fireint	kg m-2 s-1	gridcell total	daily (monthly)	
*Carbon loss due to peat burning	somcfire	kg m-2 s-1	gridcell total	monthly	
Carbon Mass Flux out of Atmosphere due to Net biome Production on Land (NBP)	ecoatmflux	kg m-2 s-1	gridcell total	daily (monthly)	This is the net mass flux of carbon between land and atmosphere calculated as photosynthesis MINUS the sum of plant and soil respiration, carbon fluxes from fire, harvest, grazing and land-use change. Positive flux is into the land.
Structure [as Biomes output Table]					
Leaf Area Index	lai-<pft>	1	per pft	annual	
Leaf Area Index	lai-<pft>	1	gridcell average	daily (monthly)	
Plant Functional Type Grid Fraction	pft-<pft>	%	per gridcell	annual (or once if static)	The categories may differ from model to model, depending on their PFT definitions. This may include natural PFTs, anthropogenic PFTs, bare soil, lakes, urban areas, etc. Sum of all should equal the fraction of the grid-cell that is land.
Hydrological variables [as per Biomes output Table]					
Runoff	qtot	kg m-2 s-1	per gridcell	daily** (monthly)	Total (surface + subsurface) runoff (qtot = qs + qsb). If daily resolution not possible, please provide monthly. If storage issues keep you from reporting daily data, please contact the ISI-MIP team to discuss potential solutions. **For those models also participating in the water simulations

Soil moisture	soilmoist	kg m ⁻²	per grid cell	monthly	Please provide soil moisture for all depth layers (i.e., 3D-field), and indicate depth in m.
Frozen soil moisture for each layer	soilmoistfroz	kg m ⁻²	per gridcell	monthly	Please provide frozen soil moisture for all depth levels and indicate depth in m.
Snow depth	snd	m	per gridcell	daily (monthly)	Grid cell mean depth of snowpack.
Snow water equivalent	swe	kg m ⁻²	per gridcell	daily (monthly)	Total water mass of the snowpack (liquid or frozen), averaged over a grid cell.
Annual maximum thaw depth	thawdepth	m	per gridcell	annual	Calculated from daily thaw depths
Other outputs					
Burnt Area Fraction	burntarea	%	per gridcell	daily (monthly)	Fraction of entire gridcell that is covered by burnt vegetation
N ₂ O emissions into atmosphere	n2o	kg m ⁻² s ⁻¹	gridcell total	monthly	From land, not from industrial fossil fuel emissions and transport
CH ₄ emissions into atmosphere	ch4	kg m ⁻² s ⁻¹	gridcell total	monthly	From land, not from industrial fossil fuel emissions and transport

15 References

- Bolt, J. and van Zanden, J. L.: The Maddison Project: collaborative research on historical national accounts, *Econ. Hist. Rev.*, 67(3), 627–651, 2014.
- Choulga, M., Kourzeneva, E., Zakharova, E. and Doganovsky, A.: Estimation of the mean depth of boreal lakes for use in numerical weather prediction and climate modelling, *Tellus A Dyn. Meteorol. Oceanogr.*, 66(1), 21295, doi:10.3402/tellusa.v66.21295, 2014.
- Dellink, R., Chateau, J., Lanzi, E. and Magné, B.: Long-term economic growth projections in the Shared Socioeconomic Pathways, *Glob. Environ. Chang.*, doi:10.1016/j.gloenvcha.2015.06.004, 2015. [¶](#)
- [Elliott, J. and Müller, C. and Deryng, D. and Chryssanthacopoulos, J. and Boote, K. J. and Büchner, M. and Foster, I. and Glotter, M. and Heinke, J. and Iizumi, T. and Izaurrealde, R. C. and Mueller, N. D. and Ray, D. K. and Rosenzweig, C. and Ruane, A. C. and Sheffield, J.: The Global Gridded Crop Model Intercomparison: data and modeling protocols for Phase 1 \(v1.0\), *Geosci. Model Dev.*, 8, 261–277, <https://doi.org/10.5194/gmd-8-261-2015>, 2015.](#)
- Frieler, K., Lange, S., Piontek, F., Reyer, C. P. O., Schewe, J., Warszawski, L., Zhao, F., Chini, L., Denvil, S., Emanuel, K., Geiger, T., Halladay, K., Hurtt, G., Mengel, M., Murakami, D., Ostberg, S., Popp, A., Riva, R., Stevanovic, M., Suzuki, T., Volkholz, J., Burke, E., Ciais, P., Ebi, K., Eddy, T. D., Elliott, J., Galbraith, E., Gosling, S. N., Hattermann, F., Hickler, T., Hinkel, J., Hof, C., Huber, V., Jägermeyr, J., Krysanova, V., Marcé, R., Müller Schmied, H., Mouratiadou, I., Pierson, D., Tittensor, D. P., Vautard, R., van Vliet, M., Biber, M. F., Betts, R. A., Bodirsky, B. L., Deryng, D., Froking, S., Jones, C. D., Lotze, H. K., Lotze-Campen, H., Sahajpal, R., Thonicke, K., Tian, H., and Yamagata, Y.: Assessing the impacts of 1.5 °C global warming – simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b), *Geosci. Model Dev.*, 10, 4321–4345, <https://doi.org/10.5194/gmd-10-4321-2017>, 2017.
- [Gasparrini A, Leone M. Attributable risk from distributed lag models. *BMC Med Res Methodol.* 2014 Apr 23;14:55. doi: 10.1186/1471-2288-14-55. PMID: 24758509; PMCID: PMC4021419. ¶](#)
- Haith, D. A. and Shoemaker, L. L.: Generalized Watershed Loading Functions for stream flow nutrients, *Water Resour. Bull.*, 23, 471–478, 1987.
- [Håkanson, L. Models to predict Secchi depth in small glacial lakes. *Aquatic Science* 57, 31–53 \(1995\). <https://doi.org/10.1007/BF00878025>¶](#)
- [Hinkel, Jochen and Lincke, Daniel and Vafeidis, Athanasios T. and Perrette, Mahé and Nicholls, Robert James and Tol, Richard S. J. and Marzeion, Ben and Fettweis, Xavier and Ionescu, Cezar and Levermann, Anders: Coastal flood damage and adaptation costs under 21st century sea-level rise, *Proceedings of the National Academy of Sciences*, 111 \(9\): 3292–3297; DOI: 10.1073/pnas.1222469111, 2014.¶](#)
- Hurtt, G. C., L. Chini, R. Sahajpal, S. Froking, B. L. Bodirsky, K. Calvin, J. C. Doelman, J. Fisk, S. Fujimori, K. K. Goldewijk, T. Hasegawa, P. Havlik, A. Heinemann, F. Humpenöder, J. Jungclaus, Jed Kaplan, J. Kennedy, T. Kristzin, D. Lawrence, P. Lawrence, L. Ma, O. Mertz, J. Pongratz, A. Popp, B. Poulter, K. Riahi, E. Shevliakova, E. Stehfest, P. Thornton, F. N. Tubiello, D. P. van Vuuren, X. Zhang (2020). Harmonization of Global Land-Use Change and Management for the Period 850–2100 (LUH2) for CMIP6. *Geoscientific Model Development Discussions*. <https://doi.org/10.5194/gmd-2019-360>
- Klein Goldewijk, K., Beusen, A., Doelman, J., and Stehfest, E.: Anthropogenic land use estimates for the Holocene – HYDE 3.2, *Earth Syst. Sci. Data*, 9, 927–953, <https://doi.org/10.5194/essd-9-927-2017>, 2017. [¶](#)
- [Kopp, Robert E. and Horton, Radley M. and Little, Christopher M. and Mitrovica, Jerry X. and Oppenheimer, Michael and Rasmussen, D. J. and Strauss, Benjamin H. and Tebaldi, Claudia: Probabilistic 21st and 22nd century sea-level projections at a global network of tide-gauge sites, *Earth's Future*, 2 \(8\): 383–406, <https://doi.org/10.1002/2014EF000239>, 2014.¶](#)

- Kopp, Robert E. and Kemp, Andrew C. and Bittermann, Klaus and Horton, Benjamin P. and Donnelly, Jeffrey P. and Gehrels, W. Roland and Hay, Carling C. and Mitrovica, Jerry X. and Morrow, Eric D. and Rahmstorf, Stefan: Temperature-driven global sea-level variability in the Common Era, Proceedings of the National Academy of Sciences, 113 (11): E1434--E1441, doi:10.1073/pnas.1517056113, 2016.¶
- Kourzeneva, E. 2009. Global dataset for the parameterization of lakes in numerical weather prediction and climate modelling. ALADIN Newsletter. 37. July December, (eds. F. Bouttier and C. Fischer), Meteo-France, Toulouse, France, 46 53.
- Kourzeneva, E.: External data for lake parameterization in Numerical Weather Prediction and climate modeling, *Boreal Environ. Res.*, 15(2), 165–177, 2010.
- Lange, S.: Bias correction of surface downwelling longwave and shortwave radiation for the EWEMBI dataset, Earth Syst. Dynam., 9, 627–645, https://doi.org/10.5194/esd-9-627-2018, 2018.¶
- Lamarque, J. F., Dentener, F., McConnell, J., Ro, C. U., Shaw, M., Vet, R., Bergmann, D., Cameron-Smith, P., Dalsoren, S., Doherty, R., Faluvegi, G., Ghan, S. J., Josse, B., Lee, Y. H., Mackenzie, I. a., Plummer, D., Shindell, D. T., Skeie, R. B., Stevenson, D. S., Strode, S., Zeng, G., Curran, M., Dahl-Jensen, D., Das, S., Fritzsche, D. and Nolan, M.: Multi-model mean nitrogen and sulfur deposition from the atmospheric chemistry and climate model intercomparison project (ACCMIP): Evaluation of historical and projected future changes, *Atmos. Chem. Phys.*, 13(16), 7997–8018, doi:10.5194/acp-13-7997-2013, 2013a.
- Lamarque, J. F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. a., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, a. and Zeng, G.: The atmospheric chemistry and climate model intercomparison Project (ACCMIP): Overview and description of models, simulations and climate diagnostics, *Geosci. Model Dev.*, 6(1), 179–206, doi:10.5194/gmd-6-179-2013, 2013b.
- De Lary, R.: Massif des Landes de Gascogne. II – ETAT DES CONNAISSANCES TECHNIQUES, Bourdeaux., 2015.
- Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296(1–4), 1–22, doi:10.1016/J.JHYDROL.2004.03.028, 2004.
- Millero FJ & Poisson A: International one-atmosphere equation of state of seawater. *Deep-Sea Research*, 28, 625–629, 1981.
- Monfreda, C., Ramankutty, N. and Foley, J. A.: Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000, *Glob. Biogeochem. Cycles*, 22(GB1022), doi:10.1029/2007GB002947., 2008.
- Müller Schmied, H., Adam, L., Eisner, S., Fink, G., Flörke, M., Kim, H., Oki, T., Portmann, F. T., Reinecke, R., Riedel, C., Song, Q., Zhang, J. and Döll, P.: Impact of climate forcing uncertainty and human water use on global and continental water balance components, *Proc. Int. Assoc. Hydrol. Sci.*, 93, doi:10.5194/piahs-93-1-2016, 2016.
- Murakami, D. and Yamagata, Y.: Estimation of gridded population and GDP scenarios with spatially explicit statistical downscaling, [online] Available from: <http://arxiv.org/abs/1610.09041> (Accessed 29 May 2017), 2016.
- Popp, A., Humpenöder, F., Weindl, I., Bodirsky, B. L., Bonsch, M., Lotze-Campen, H., Müller, C., Biewald, A., Rolinski, S., Stevanovic, M. and Dietrich, J. P.: Land-use protection for climate change mitigation, *Nat. Clim. Chang.*, 4(December), 2–5, doi:10.1038/nclimate2444, 2014.

[Reyer, C. P. O., Silveyra Gonzalez, R., Dolos, K., Hartig, F., Hauf, Y., Noack, M., Lasch-Born, P., Rötzer, T., Pretzsch, H., Meesenburg, H., Fleck, S., Wagner, M., Bolte, A., Sanders, T. G. M., Kolari, P., Mäkelä, A., Vesala, T., Mammarella, I., Pumpanen, J., Collalti, A., Trotta, C., Matteucci, G., D'Andrea, E., Foltýnová, L., Krejza, J., Ibrom, A., Pilegaard, K., Loustau, D., Bonnefond, J.-M., Berbigier, P., Picart, D., Lafont, S., Dietze, M., Cameron, D., Vieno, M., Tian, H., Palacios-Orueta, A., Cicuendez, V., Recuero, L., Wiese, K., Büchner, M., Lange, S., Volkholz, J., Kim, H., Horemans, J. A., Bohn, F., Steinkamp, J., Chikalanov, A., Weedon, G. P., Sheffield, J., Babst, F., Vega del Valle, I., Suckow, F., Martel, S., Mahnken, M., Gutsch, M., and Frieler, K.: The PROFOUND Database for evaluating vegetation models and simulating climate impacts on European forests, *Earth Syst. Sci. Data*, 12, 1295–1320, <https://doi.org/10.5194/essd-12-1295-2020>, 2020. ¶](#)

Samir, C. and Lutz, W.: The human core of the shared socioeconomic pathways: Population scenarios by age, sex and level of education for all countries to 2100, *Glob. Environ. Chang.*, doi:10.1016/j.gloenvcha.2014.06.004, 2014.

Schneiderman, E. M., Pierson, D. C., Lounsbury, D. G. and Zion, M. S.: Modeling the hydrochemistry of the Cannonsville watershed with Generalized Watershed Loading Functions (GWLF), *J. Am. Water Resour. Assoc.*, 38, 1323–1347, 2002. ¶

[Shatwell \(unpubl.\)](#)

Stevanović, M., Popp, A., Lotze-Campen, H., Dietrich, J. P., Müller, C., Bonsch, M., Schmitz, C., Bodirsky, B., Humpenöder, F. and Weindl, I.: High-end climate change impacts on agricultural welfare, *Sci. Adv.*, 2016.

Subin, Z. M., Riley, W. J. and Mironov, D.: An improved lake model for climate simulations: Model structure, evaluation, and sensitivity analyses in CESM1, *J. Adv. Model. Earth Syst.*, 4(1), M02001, doi:10.1029/2011MS000072, 2012.

Wada, Y., Flörke, M., Hanasaki, N., Eisner, S., Fischer, G., Tramberend, S., Satoh, Y., Van Vliet, M. T. H., Yillia, P., Ringler, C., Burek, P. and Wiberg, D.: Modeling global water use for the 21st century: The Water Futures and Solutions (WFaS) initiative and its approaches, *Geosci. Model Dev.*, 9(1), 175–222, doi:10.5194/gmd-9-175-2016, 2016. ¶